

GeDiPNet Shared-Gene Disease Pairs

Similarity: Jaccard index on shared curated genes (disease_gdp). Significance: upper-tail hypergeometric test ($P(X \geq k)$), Benjamini-Hochberg FDR-corrected across all tested pairs. Corroborated Shared Genes: count of shared genes backed by ≥ 2 distinct disease_gdp.main_source values combined across both diseases (e.g. GWAS-backed for one, ClinVar-backed for the other still counts); Shared Gene Names lists each gene's own source count in parentheses. Overlap Coefficient: shared genes / MIN(each disease's own total gene count) -- complements the Jaccard-based Similarity Score by showing how much of the SMALLER disease's entire known gene-set is covered by this pair, which Jaccard (dividing by the union) can understate. Shared Cluster: cluster id if both diseases fall in the same label-propagation cluster (scripts/build_disease_clusters.php), blank otherwise. See scripts/build_related.php on gedipnet.bicnirrh.res.in. Data as of: 2026-10-09 03:00:13. Filters applied: disease name contains "Congenital ectodermal dysplasia of face" (5 matching pairs). · Rows: 5 · Generated: 2026-10-09 20:06:06

Disease A	Disease B	Shared Genes	Corroborated Shared Genes	Shared Gene Names	Similarity Score	Overlap Coefficient	P-value	FDR Q-value	Shared Cluster
Ablepharon macrostomia syndrome	Congenital ectodermal dysplasia of face	1	0	TWIST2 (1)	0.50000	1.00000	6.494e-5	2.329e-4	
Barber-say syndrome	Congenital ectodermal dysplasia of face	1	1	TWIST2 (6)	0.50000	1.00000	6.494e-5	2.329e-4	
Congenital ectodermal dysplasia of face	Focal facial dermal dysplasia	1	1	TWIST2 (5)	0.33333	1.00000	1.299e-4	3.929e-4	
Congenital ectodermal dysplasia of face	Dry eye syndrome	1	1	TWIST2 (2)	0.01724	1.00000	3.702e-3	4.734e-3	
Congenital ectodermal dysplasia of face	Sjogren syndrome	1	1	TWIST2 (2)	0.01389	1.00000	4.611e-3	5.731e-3	