

GeDiPNet Shared-Gene Disease Pairs

Similarity: Jaccard index on shared curated genes (disease_gdp). Significance: upper-tail hypergeometric test ($P(X \geq k)$), Benjamini-Hochberg FDR-corrected across all tested pairs. Corroborated Shared Genes: count of shared genes backed by ≥ 2 distinct disease_gdp.main_source values combined across both diseases (e.g. GWAS-backed for one, ClinVar-backed for the other still counts); Shared Gene Names lists each gene's own source count in parentheses. Overlap Coefficient: shared genes / MIN(each disease's own total gene count) -- complements the Jaccard-based Similarity Score by showing how much of the SMALLER disease's entire known gene-set is covered by this pair, which Jaccard (dividing by the union) can understate. Shared Cluster: cluster id if both diseases fall in the same label-propagation cluster (scripts/build_disease_clusters.php), blank otherwise. See scripts/build_related.php on gedipnet.bicnirrh.res.in. Data as of: 2026-10-21 12:58:11. Filters applied: disease name contains "Hypoplasia or aplasia of tibia with polydactyly" (5 matching pairs). · Rows: 5 · Generated: 2026-10-21 16:18:53

Disease A	Disease B	Shared Genes	Corroborated Shared Genes	Shared Gene Names	Similarity Score	Overlap Coefficient	P-value	FDR Q-value	Shared Cluster
Hypoplasia or aplasia of tibia with polydactyly	Triphalangeal thumb-polysyndactyly syndrome	2	2	LMBR1 (4), SHH (3)	0.66667	1.00000	8.338e-9	6.594e-8	
Hypoplasia or aplasia of tibia with polydactyly	Laurin-sandrow syndrome	2	0	LMBR1 (1), SHH (1)	0.66667	1.00000	8.338e-9	6.594e-8	
Hypoplasia or aplasia of tibia with polydactyly	Preaxial polydactyly 2	2	0	LMBR1 (1), SHH (1)	0.66667	1.00000	8.338e-9	6.594e-8	
Acheiropody	Hypoplasia or aplasia of tibia with polydactyly	1	1	LMBR1 (2)	0.33333	1.00000	1.291e-4	3.269e-4	
Hypoplasia or aplasia of tibia with polydactyly	Patterson stevenson fontaine syndrome	1	1	LMBR1 (2)	0.33333	1.00000	1.291e-4	3.269e-4	